



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROOF OF A FORMULA DUE TO CAUCHY.

By PROF. W. H. ECHOLS, Charlottesville, Va.

The function

$$\begin{vmatrix} 1 & , & 1 & , & 1 & , & 1 & , & \dots & , & 1 \\ a^p x & , & 1 & , & a & , & a^2 & , & \dots & , & a^{n-1} \\ (a^p x)^2 & , & 1 & , & a^2 & , & (a^2)^2 & , & \dots & , & (a^{n-1})^2 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ (a^p x)^n & , & 1 & , & a^n & , & (a^2)^n & , & \dots & , & (a^{n-1})^n \end{vmatrix} \div \zeta^{\frac{1}{2}}(a, a^2 \dots a^n), \quad (1)$$

vanishes when $x = a^{i-p}$ ($i = 0, 1, \dots, n-1$), p being any finite quantity.

If we expand (1) with respect to the first column, we have

$$1 + \sum_{r=1}^{r=\infty} (-1)^r \frac{(1 - a^{-n+r-1}) \dots (1 - a^{-n})}{(a-1) \dots (a^r-1)} a^{r(p+1)} x^r.$$

This rational integral function of x of the n th degree has the n roots a^{-p} , a^{1-p} , \dots , a^{n-1-p} ; whence we have the identity

$$\prod_{i=0}^{i=n-1} (1 - a^{p-i} x) = 1 + \sum_{r=1}^{r=\infty} (-1)^r \frac{(1 - a^{-n+r-1}) \dots (1 - a^{-n})}{(a-1) \dots (a^r-1)} a^{r(p+1)} x^r, \quad (2)$$

Evidently, if $a > 1$, we have

$$\prod_{i=0}^{i=\infty} (1 - a^{p-i} x) = 1 + \sum_{r=1}^{r=\infty} \frac{(-1)^r a^{r(p+1)}}{(a-1) \dots (a^r-1)} x^r. \quad (3)$$

Let $x = a^{q-p}$, where q is any positive integer or zero, then

$$0 = 1 + \sum_{r=1}^{r=\infty} \frac{(-1)^r a^{r(q+1)}}{(a-1) \dots (a^r-1)}.$$

In (2) let $p = -1$, we then have Cauchy's formula (Comptes Rendus, 1840. Chrystal's Algebra, II, 316), and (3) becomes Euler's theorem (Introd. in Anal. Inf., § 306).

As a consequence the sum of the products, taken q at a time without repetition, of the quantities

$$a^{-i} \quad (i = 1, 2, 3, \dots, \infty)$$

is

$$\frac{1}{(a-1) \dots (a^q-1)}.$$